



## Atorvastatin Drug Product Analyzed with HPLC - AppNote

### API well separated from degradation product

Using this Simple Linear Gradient, Atorvastatin is well separated from its main degradation Product. The degradant is formed under acidic conditions and therefore its Separation is important for Stability Indicating Methods for pharmaceutical formulations of Atorvastatin. For Methods requiring non-degrading conditions, it is important to choose diluent and Mobile Phase conditions with pH sensitivity in mind.

As such, Ammonium Acetate was chosen for the diluent and Mobile Phase additives in order to avoid any degradation from these sources. Data obtained under non-degrading conditions is shown in *Figure A*. An overlay of five consecutive runs of the degraded extract is shown in *Figure B*. The Repeatability of the Method is demonstrated by the consistent retention time (%RSD of 0.10).





### Peaks:

1. Atorvastatin
2. Degradant

### Method Conditions

**Column:** Cogent Bidentate C18™, 4µm, 100Å

**Catalog No.:** [40018-15P](#)

**Dimensions:** 4.6 x 150mm

#### Mobile Phase:

A: DI Water / 10mM Ammonium Acetate

B: 90% Acetonitrile / 10% DI Water / 10mM Ammonium Acetate

#### Gradient:

Time (minutes)	%B
0	40
5	100
6	40

**Injection vol.:** 2µL

**Flow rate:** 0.8mL / minute

**Detection:** UV @ 248nm

#### Sample Preparation:

**A:** Atorvastatin tablet extract

**B:** Degraded Atorvastatin tablet extract

40mg strength tablet ground was placed in 100mL volumetric flask. 50mL of Mobile phase B was added to flask. Sample was vortexed for 5 minutes, sonicated for 5 minutes, then diluted to mark with Solution A. A portion was filtered through a 0.45µm Nylon Syringe Filter (MICROSOLV Tech Corp.). Final Working Solution was then diluted 4x in **Sample A**) 50:50 A / B or **Sample B**) 1.0 N HCL in 50:50 A / B.

---

**Note:** Atorvastatin is a competitive inhibitor of 3-Hydroxy-3-Methylglutaryl Co-enzyme A (HMGCoA) reductase, which catalyzes the rate-limiting step in Cholesterol biosynthesis. As such, A torvastatin is used to reduce plasma levels of low-density lipoprotein (LDL) Cholesterol, which are known to contribute to the development of Atherosclerosis.

---



**Attachment Atorvastatin Drug Product Analyzed with HPLC pdf [Download File](#)**

Printed from the Chrom Resource Center

Copyright 2025, All Rights Apply

**MicroSolv Technology Corporation**

9158 Industrial Blvd. NE, Leland, NC 28451

Tel: (732) 380-8900

Fax: (910) 769-9435

Email: [customers@mtc-usa.com](mailto:customers@mtc-usa.com)

Website: [www.mtc-usa.com](http://www.mtc-usa.com)