



Furazolidone Analyzed with HPLC - AppNote

API and Related Compounds Separation

Furazolidone was Separated from related compounds in a commercially available pharmaceutical preparation. Excellent Peak shape and Selectivity from the related compounds was achieved with this Robust Method making it a great choice for QC applications even when using aggressive Mobile Phases that include acids. Molecular weight of Furazolidone is 225.16.



Peak:

Furazolidone

Method Conditions

Column: Cogent Bidentate C18™, 4 µm, 100 Å

Catalog No.: [40018-15P](#)

Dimensions: 4.6 x 150 mm

Mobile Phase: 12% Acetonitrile / 88% DI Water / 1% Acetic Acid

Temperature: 30°C

Injection vol.: 20 µL

Flow rate: 2mL / minute

Detection: UV @ 365 nm

Note: *This antibiotic drug is used to treat diarrhea and enteritis caused by bacteria or protozoan infection. Also used as a veterinary medicine for salmonids and Myxobolus cerebralis infection.*



Attachment No 73 Furazolidone Analyzed with HPLC pdf 0.2 Mb [Download File](#)

Printed from the Chrom Resource Center

Copyright 2025, All Rights Apply

MicroSolv Technology Corporation

9158 Industrial Blvd. NE, Leland, NC 28451

Tel: (732) 380-8900

Fax: (910) 769-9435

Email: customers@mtc-usa.com

Website: www.mtc-usa.com