



Isomers of Ascorbic Acid D and L Separation by HPLC - AppNote

An Analysis of Vitamin C

Click [HERE](#) for Column Ordering Information.

This simple Method shows good separation between two Enantiomers as well as Retention, considering these Acids both have a - 1.6 log p. This Method is Robust as results were verified with three different HPLCs and two separate Columns, all showing excellent Resolution of these two Acids.



Peaks:

1. D- Isoascorbic acid
2. L- Ascorbic acid

Method Conditions

Column: Cogent Diamond Hydride™, 4µm, 100Å

Catalog No.: [70000-10P](#)

Dimensions: 4.6mm x 100mm

Mobile Phase: 98% Acetonitrile 2% DI Water / 0.1% Formic Acid

Flow Rate: 1.0mL / minute

Injection Volume: 1uL

Detection: UV 254nm

Injection vol.: 1µL

Sample Preparation: D- Isoascorbic Acid and L- Ascorbic Acid in 1.0 mg/mL in diluent of 50% Acetonitrile / 50% DI Water (v/v)

t₀: 1.20 Minutes

K₁: 0.39

K₂: 0.62

α: 1.59

Note: Ascorbic acid exists as two enantiomers (mirror-image isomers), commonly denoted "l" (for "levo") and "d" (for "dextro"). The l isomer is the one most often encountered and occurs naturally in many foods, and is one form of Vitamin C, an essential nutrient for many animals. Deficiency of Vitamin C causes scurvy. Vitamin C is used as a food additive and a dietary supplement for its antioxidant properties. The "d" form can be made via chemical synthesis but has no significant biological role.

Capacity Factor - Relative Retention $k = (t_R - t_0) / t_0$

$\alpha = K_2 / K_1$



Printed from the Chrom Resource Center

Copyright 2025, All Rights Apply

MicroSolv Technology Corporation

9158 Industrial Blvd. NE, Leland, NC 28451

Tel: (732) 380-8900

Fax: (910) 769-9435

Email: customers@mtc-usa.com

Website: www.mtc-usa.com