





Atenolol Tablet Analyzed with HPLC - AppNote

Quick Analysis without Ion Pair Agents

The USP Assay Method for Atenolol uses Heptane Sulfonate and Dibutyl Amine in the Mobile Phase, which are slow to load and remove from a Reversed Phase HPLC Column. These reagents are added to improve the Peak Tailing, which must be not more than 2.0.

The Method in this Application Note is an improvement over the official USP Method as the Peak produced is very Symmetrical without the use of Ion Pair Agents. Also, the Method is LCMS compatible if transfer is desired. Data from two Column lots is shown to illustrate lot-to-lot consistency of this Method.



Peak:

Atenolol

Method Conditions

Column: Cogent Diamond Hydride™, 4µm, 100Å

Catalog No.: 70000-7.5P

Dimensions: 4.6 x 75mm

Mobile Phase:

A: DI Water with 0.1% Formic Acid (v/v)

B: Acetonitrile with 0.1% Formic Acid (v/v)

Gradient:

Time (minutes)	%B
0	90
1	90
6	40
7	90

Post Time: 3 minutes

Injection vol.: 2µL

Flow rate: 1.0mL / minute

Detection: UV @ 225nm

Sample Preparation: 25mg strength Atenolol tablet was ground and added to a 25mL volumetric flask. A portion of 50:50 Solvent A / Solvent B diluent was added and the flask was sonicated 10 minutes. It was then diluted to mark and mixed. A portion was filtered with a 0.45µm Nylon Syringe Filter (MICROSOLV Tech Corp.) and diluted 1:10.

t₀: 0.9 minutes

Note : Atenolol is a selective β 1 receptor antagonist. It is used to treat various cardiovascular diseases such as hypertension. It is available by prescription under the trade name Tenormin®.



Attachment No 250 Atenolol Tablet Analyzed with HPLC pdf 0.4 Mb [Download File](#)

Printed from the Chrom Resource Center

Tel: (732) 380-8900

Copyright 2025, All Rights Apply

Fax: (910) 769-9435

MicroSolv Technology Corporation

Email: customers@mtc-usa.com

9158 Industrial Blvd. NE, Leland, NC 28451

Website: www.mtc-usa.com