


# Lidocaine in Gel Analyzed with HPLC - AppNote

# **Assay Method for Solarcaine® Gel Extract**

This AppNote shows how Lidocaine Hydrochloride (HCL) from a gel extract can be easily analyzed. The compound has a tertiary amine but excellent Peak shape is achieved; which is often difficult to obtain for analytes like this.

A Gradient was used to ensure components from the gel extract matrix do not build up on the Column and adversely affect robustness or Column life. The Column robustness is demonstrated by the overlay of injections from two different Column lots.



#### Peak:

1. Lidocaine

### **Method Conditions**

Column: Cogent Diamond Hydride™, 4µm, 100Å

Catalog No.: 70000-7.5P Dimensions: 4.6 x 75mm

Mobile Phase:

A: DI Water / 0.1% Trifluoroacetic Acid (v/v) B: Acetonitrile / 0.1% Trifluoroacetic Acid (v/v)

Gradient:

Time (minutes)	%B
0	97
1.5	97
6	40
7	97

Post Time: 3 minutes Injection vol.: 2µL

Flow rate: 1.0mL / minutes Detection: UV @ 220nm

**Sample Preparation**: 250mg Solarcaine gel containing 0.5% Lidocaine HCL was weighed in a 25mL volumetric flask. A portion of 50:50 Solvent A / Solvent B diluent was added and the flask was sonicated 15 minutes at 40°C. It was then diluted to mark and filtered with a 0.45µm Nylon Syringe Filter (MICROSOLV Tech Corp.).

**to**: 0.9 minutes

Note: Lidocaine is a local anesthetic used in a variety of applications such as sunburn relief and numbing for dental procedures. As such, several types of formulations are used such as topical creams / gels and injectable solutions. Trade names include Solarcaine and Xylocaine®.



## **Attachment**

No 215 Lidocaine in Gel Analyzed with HPLC pdf 0.4 Mb Download File

Printed from the Chrom Resource Center
Copyright 2025, All Rights Apply
MicroSolv Technology Corporation
9158 Industrial Blvd. NE, Leland, NC 28451

Tel: (732) 380-8900
Fax: (910) 769-9435
Email: customers@mtc-usa.com
Website: www.mtc-usa.com