



Cytidine Based Cancer Prodrug Analyzed with HPLC – AppNote

Separation Achieved with Shape Selectivity and Hydrophobic Interaction

This Method provides an example of Reversed Phase Separation containing a number of related compounds. The properties of the solutes are listed in the Table below.

When comparing the properties of Compounds 2 and 3, it is interesting to note that 2 is more hydrophobic than 3 but its Retention is shorter. The Method uses Shape Selectivity as well as Hydrophobic Interaction which may explain why the less hydrophobic compound is retained longer. In general, the more polar compounds elute earlier than the more hydrophobic species. Compound 7 has no polar functionalities at all so it is the most retained under Reversed Phase conditions.



Peaks:

1. Cytidine-R1
2. Cytidine-R2
3. Cytidine-R3
4. Quinolinedione-R1
5. Saccharin
6. Quinolinedione-R2
7. Benzopyran

Method Conditions

Column: Cogent UDC Cholesterol™, 4 µm, 100Å

Catalog No.: 69069-7.5P

Dimensions: 4.6 x 75 mm

Mobile Phase: 60:40 Acetonitrile / DI Water + 0.5% Perchloric Acid

Injection vol.: 5 µL

Flow rate: 1 mL / minute

Detection: UV @ 270nm

Sample Preparation: 1 mg / mL of each in Acetonitrile + DI Water

Table:

Solute	Mol Wt.	B pKa	Log P
Cytidine-R1	397	3.73	1.54
Cytidine-R2	454	3.73	3.66
Cytidine-R3	425	3.73	2.66
Quinolinedione-R1	536	-	3.46
Saccharin	536	5.36	2.96
Quinolinedione-R2	520	-	2.74
Benzopyran	396	-	2.80



Attachment Cancer Prodrug Analyzed with HPLC pdf 0.2 Mb [Download File](#)

Printed from the Chrom Resource Center

Copyright 2025, All Rights Apply

MicroSolv Technology Corporation

9158 Industrial Blvd. NE, Leland, NC 28451

Tel: (732) 380-8900

Fax: (910) 769-9435

Email: customers@mtc-usa.com

Website: www.mtc-usa.com